

## Aberystwyth University

### *Microsatellite analysis supports the existence of three cryptic species within the bumble bee *Bombus lucorum sensu lato**

McKendrick, Lorraine; Provan, James; Fitzpatrick, Úna; Brown, Mark J. F.; Murray, Tomás E.; Stolle, Eckart; Paxton, Robert J.

*Published in:*  
Conservation Genetics

*DOI:*  
[10.1007/s10592-017-0965-3](https://doi.org/10.1007/s10592-017-0965-3)

*Publication date:*  
2017

*Citation for published version (APA):*

McKendrick, L., Provan, J., Fitzpatrick, Ú., Brown, M. J. F., Murray, T. E., Stolle, E., & Paxton, R. J. (2017). Microsatellite analysis supports the existence of three cryptic species within the bumble bee *Bombus lucorum sensu lato*. *Conservation Genetics*, 18(3), 573-584. <https://doi.org/10.1007/s10592-017-0965-3>

#### General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)

# **Microsatellite analysis supports the existence of three cryptic species within the bumble bee *Bombus lucorum sensu lato***

Lorraine McKendrick<sup>1</sup>, Jim Provan<sup>2</sup>, Úna Fitzpatrick<sup>3</sup>, Mark J. F. Brown<sup>4</sup>, Tomás E. Murray<sup>3</sup>,  
Eckart Stolle<sup>5</sup>, Robert J. Paxton<sup>1,6 \*</sup>

<sup>1</sup> School of Biological Sciences, Queen's University Belfast, Belfast BT9 7BL, UK

<sup>2</sup> Institute of Biological, Environmental and Rural Sciences, Penglais, Aberystwyth University, Aberystwyth SY23 3DA, UK

<sup>3</sup> National Biodiversity Data Centre, Carriganore, Waterford, Ireland

<sup>4</sup> School of Biological Sciences, Royal Holloway University of London, Egham, Surrey TW20 0EX, UK

<sup>5</sup> School of Biological and Chemical Sciences, Queen Mary University of London, Mile End Road, London E1 4NS, UK

<sup>6</sup> Institute for Biology, Martin Luther-University Halle-Wittenberg, Hoher Weg 8, 06120 Halle (Saale), Germany and German Centre for Integrative Biodiversity Research Halle-Jena-Leipzig (iDiv), Deutscher Platz 5e, 04103 Leipzig, Germany

\*Corresponding author: Robert Paxton

E-mail: [robert.paxton@zoologie.uni-halle.de](mailto:robert.paxton@zoologie.uni-halle.de)

Tel: +49 (0)345 55 26500

Fax: +49 (0)345 55 27428

## Abstract

Mitochondrial cytochrome oxidase I (COI) partial sequences are widely used in taxonomy for species identification. Increasingly, these sequence identities are combined with modeling approaches to delineate species. Yet the validity of species delineation based on such DNA ‘barcodes’ is rarely tested and may be called into question by phenomena such as ancestral polymorphisms in DNA sequences, phylogeographic divergence, mitochondrial introgression and hybridization, or distortion of mitochondrial inheritance through such factors as *Wolbachia* infection. The common and widespread European bumble bee *Bombus lucorum* s. lato contains three distinct mitochondrial DNA lineages that are assumed to represent three cryptic species, namely *Bombus cryptarum*, *B. lucorum* s. str. and *Bombus magnus*. To test whether nuclear gene pools of the three putative species were differentiated, we genotyped 304 sympatric members of the *lucorum* complex (54 *B. cryptarum* females, 168 *B. lucorum* s. str. females and 82 *B. magnus* females, as defined using mtDNA COI haplotypes) from 11 localities spread across the island of Ireland at seven nuclear microsatellite loci. Multilocus genotypes clustered into three discrete groups that largely corresponded to the three mtDNA lineages: *B. cryptarum*, *B. lucorum* s. str. and *B. magnus*. The good fit of mitochondrial haplotype to nuclear (microsatellite) genotypic data supports the view that these three bumble bee taxa are reproductively isolated species, as well as providing a vindication of species identity using so-called DNA barcodes.

**Keywords** DNA barcode; *cryptarum*; *magnus*; mitochondrial cytochrome oxidase I; STRUCTURE software; PCoA; DAPC, sympatry

## Introduction

Bumble bees (Hymenoptera: Apidae, genus *Bombus*) are of great ecological and economic importance as major pollinators of both crops and wild flowers in the Northern Hemisphere, yet they are in decline (e.g. Fitzpatrick et al. 2007; Goulson 2009; Cameron et al. 2011). Though members of the subgenus *Bombus sensu stricto* (= *Terrestribombus* Vogt) are the most abundant and widespread of all bumble bees, exhibiting a holarctic distribution (Hines 2008; Williams et al. 2008, 2012a), they can be difficult to identify in the field using the minor morphological differences that separate species (Carolan et al. 2012; Bossert 2015); the apparent abundance of members of the subgenus may mask the rarity of its morphologically indistinguishable, or cryptic, species.

In Europe, five species of *Bombus s. str.* are recognised: *B. cryptarum* (Fabricius), *B. lucorum* (L.), *B. magnus* Vogt, *B. terrestris* (L.) and *B. sporadicus* (Nylander). The taxonomic status of *B. terrestris* and *B. sporadicus* is widely accepted (Williams 1998). Difficulties arise over the other three species: *B. cryptarum*, *B. lucorum s. str.* and *B. magnus*, which are generally grouped to form the *lucorum* complex (*B. lucorum sensu lato*). They are cryptic species that appear very similar in colour and form, particularly as workers or males, and that are often difficult to differentiate morphologically from *B. terrestris* even as queens (Figure 1; Rasmont 1984; Rasmont et al. 1986).

Species classification based on morphological characters may not be suitable for cryptic species and genetic methods may help support species identification. Correct identification is of conservation importance because the taxonomic status of a species must be accurately established in order to assign status and direct conservation efforts (Ryder, 1986; Crandall et

al. 2000). The presence of cryptic species, however, has potentially detrimental implications since reproductively isolated groups should be managed independently of each other (Riddle et al. 2000; Palsbøll et al. 2007). To facilitate identification of cryptic species, molecular methods such as DNA barcoding can be used as a means of designating species on the basis of sequence similarity (Hebert et al. 2003). Such approaches have confirmed the view that cryptic species are particularly common in insects (e.g. Berkov 2002; Hebert et al. 2004).

In pre-DNA based studies, allozyme polymorphisms and variation in male cephalic odour bouquet supported the view that the *lucorum* complex of bumble bees comprise two or three species (reviewed in Bossert 2015). Bertsch et al. (2005) subsequently used mitochondrial cytochrome oxidase I (COI) gene sequences of specimens morphologically well-characterised as queens to show that the *lucorum* complex of bumble bees contained three distinct mitochondrial DNA (mtDNA) lineages in Europe, albeit sampling of two putative species was limited to two specimens apiece at each of two sites in Europe. Using a far larger number of samples from across Europe, including >300 from the island of Ireland, Murray et al. (2008) showed that the three mtDNA lineages exhibited considerable interspecific DNA sequence divergence ( $\geq 2.3\%$ ) at COI compared to intra-taxon sequence variability ( $\leq 1.3\%$ ), with overwhelming support for each lineage, supporting the idea that the three mtDNA lineages represent species: *B. cryptarum*, *B. lucorum s. str.* and *B. magnus*. Williams et al. (2012a) gave the distribution of these three taxa in Europe, Asia and, for *B. cryptarum*, even into North America based on COI sequences, again demonstrating overwhelming statistical support for each of the three COI lineages representing the *lucorum* complex. Murray et al. (2008) also developed a relatively quick and economic restriction enzyme based mtDNA COI marker system based on restriction fragment length polymorphisms (RFLPs) that differentiated among lineages.

Notwithstanding the success of DNA barcoding in separating species and even entire regional bee faunas (Sheffield et al. 2009; Magnacca and Brown 2010a, 2012; Schmidt et al. 2015), mitochondrial lineages may not represent independent, reproductively isolated species. Reasons include the retention of ancestral mitochondrial sequence polymorphisms, mitochondrial introgression and biased inheritance of maternal genetic markers by such factors as *Wolbachia* infection, known to be widespread in bees (Gerth et al. 2011, 2013, 2014; Gerth and Bleidorn 2013; but see Stahlhut et al. 2012), heteroplasmy (e.g., Magnacca and Brown 2010a), and associated tissue segregation of haplotypes (Magnacca and Brown 2010b). Moreover, the use of DNA barcoding or any other mitochondrial DNA marker system does not permit the detection of hybrids between taxa. This is all the more relevant for the *lucorum* complex of bumble bees, in which Carolan et al. (2012) found apparent mismatch between widely employed species-characteristic, discriminatory morphological traits of queens and mtDNA lineage among Irish specimens.

A resolution to this problem is to incorporate multilocus sequence typing into DNA barcoding studies (Gerth and Bleidorn 2013; Bossert 2015), an approach we report here for the *lucorum* complex. We used microsatellites, biparentally inherited nuclear markers, to determine the extent to which the nuclear gene pools of the *lucorum* complex taxa concur with the three mtDNA lineages of Bertsch et al. (2005) and Murray et al. (2008), with the aim of reducing taxonomic uncertainty in this group.

## Materials and Methods

### Sample collection and DNA extraction

Females (queens and workers) of *B. lucorum s.l.* were collected in 2005 and 2006 from 11 localities spread across the island of Ireland from both rural and urban environments while foraging on flowers (Table 1, Figure 2); they are a subset of the same Irish dataset originally presented in Murray et al. (2008). Individuals were either frozen or stored in 99% ethanol at 4°C prior to DNA extraction from a single leg using 10% Chelex (Walsh et al. 1991) or from half a thorax using a standard high salt protocol (Paxton et al. 1996).

#### Microsatellite genotyping and species identification by mitochondrial haplotyping

Individuals were genotyped at seven nuclear microsatellite loci (Supplementary Table S1) described in Stolle et al. (2011) and developed for *B. terrestris*. Forward primers included a 19 bp M13 5' tail (CACGACGTTGTAAAACGAC) and reverse primers included a 7 bp 5' tail (GTGTCTT). PCRs were carried out in a total volume of 10 µL containing 1-10 ng genomic DNA, 1 µM of 6-FAM-, TET- or HEX-labelled M13 primer (see Supplementary Table S1), 0.1 µM tailed forward primer, 1 µM reverse primer, 1x PCR reaction buffer (Promega), 200 µM each dNTP, 2.5 mM MgCl<sub>2</sub> and 0.25 U GoTaq Flexi DNA polymerase (Promega). PCRs were carried out on a MWG Primus thermal cycler using the following parameters: initial denaturation at 96 °C for 3 min followed by 35 cycles of denaturation at 96 °C for 45 s, annealing at 57 °C for 45 s, extension at 72 °C for 45 s, and a final extension at 72 °C for 3 min. Genotyping was carried out on an AB3730xl capillary genotyping system (Life Technologies; Carlsbad, California, USA). Allele sizes were scored using 500 LIZ size standards and were checked by comparison with previously sized samples.

Microsatellite genotypes were obtained at 5-7 loci for 304 females (54 *B. cryptarum*, 168 *B. lucorum s. str.* and 82 *B. magnus*; Table 1). All had already been classified to mt haplotype by RFLP analysis of a mitochondrial partial COI gene sequence by Murray et al. (2008), the results of which we use (and update by Sanger sequencing of the COI ‘barcode’ of eight samples) here.

## Data analysis

GENEPOP (version 3.4; Raymond and Rousset 1995) was used to test for linkage disequilibrium between nuclear microsatellite loci and for deviation from Hardy-Weinberg equilibrium (HWE) at these loci. We also tested for the presence of null alleles using MICRO-CHECKER (Van Oosterhout et al. 2004). Genetic differentiation within each putative species at microsatellite loci was calculated in MICROSATELLITE ANALYSER (MSA, version 4.05 for OSX) (Dieringer and Schlötterer 1997) and isolation by distance tested using the online web service IBDWS version 3.23 (Jensen et al. 2005).

We used three approaches to determine the fit between mtDNA lineage and multilocus nuclear genotype. In the first approach, genetic clustering of individuals was assessed using a Bayesian procedure implemented in the STRUCTURE software package (version 2.3.3; Pritchard et al. 2000). The program was run without priors, and with or without the admixture ancestry model. Twenty independent runs were carried out for each model and value of  $K$ , the number of genetic clusters, from  $K = 1$  to  $K = 3$ . Our rationale was to test the hypothesis of  $K = 3$  clusters (representing the three species: *B. cryptarum*, *B. lucorum s. str.* and *B. magnus*) versus a null hypothesis of  $K = 1$  or  $K = 2$  clusters (species). Each Markov chain Monte Carlo analysis used a burn-in of 50,000 followed by a further 500,000 iterations. STRUCTURE’s  $Q$



value, a probability of group membership, was calculated for each individual at  $K = 3$  using the admixture ancestry model.

Because our dataset suggested deviation from HWE (see results) whereas HWE is an assumption of STRUCTURE, we employed two distance-based methods to test for the association between genotypes and mitochondrial haplotypes, methods that do not make assumptions about mating structure and that do not make *a priori* assumptions about group membership. In the second approach, we visualised relationships among multilocus microsatellite genotypes of the 304 females of the *lucorum* complex using principle coordinate analysis (PCoA) in GENALEX version 6.5 (Peakall and Smouse 2006). In the third approach, we used Discriminant Analysis of Principal Components (DAPC; Jombart et al. 2010) to cluster genotypes independently of *a priori* haplotype designation using the R package *adeigenet* version 1.4.2 (Jombart 2008) in R version 3.1.0 (R Core Team 2014). For DAPC, we examined results after extracting 5, 10, 20 or 40 principal components from the genotype data.

DNA sequencing to improve mt RFLP-based haplotyping

STRUCTURE  $Q$  values suggested that three individuals were in a different genotypic cluster to that of the other individuals with the same mt RFLP haplotype (Supplementary Table S2). Preliminary visualisation of the PCoA suggested that two of these three individuals and five additional individuals were in a different genotypic cluster to those with the same mt RFLP haplotype (Supplementary Table S2). All eight aberrant individuals were sequenced at the COI 'barcode' (Hebert et al. 2003) and identified by a web-based BLASTn search against the entire NCBI nucleotide database.

The original (in Murray et al. 2008) mitochondrial RFLP classification for four of these eight individuals was incorrect; two individuals with *lucorum* RFLP patterns had *cryptarum* COI DNA sequences and two individuals with *cryptarum* RFLP patterns had *magnus* COI DNA sequences (Supplementary Table S2). Though error rates in defining an individual's mt lineage using RFLPs were likely low, they nevertheless call into question the value of using RFLPs to define unambiguously the mt haplotype, as has been proposed for the *lucorum* complex of bumble bees (Murray et al. 2008; Versterlund et al. 2014). We suggest that DNA sequencing of the COI barcode is a more reliable method of defining the mt lineage in the *lucorum* complex of bumble bees in Europe and likely in other taxa, too. We recommend Sanger sequencing rather than RFLP-based inference of haplotypes in future studies of the *lucorum* complex.

The final, updated dataset is presented in Table 1 and in the Results section below.

## Results

Approximately 10% of samples were duplicated per 96-well plate for PCR; duplicates gave identical microsatellite genotypes, suggesting very low rates of error in amplifying and calling genotypes. No consistent linkage disequilibrium (i.e. involving the same loci) was detected between any of the seven nuclear microsatellites analysed across the three putative species of the *lucorum* complex (Supplementary Table S3).

When individuals from all 11 populations were lumped together into their three putative species, *B. cryptarum*, *B. lucorum s. str.* and *B. magnus*, loci 255 and 278 exhibited deviation

from Hardy-Weinberg equilibrium in all three putative species and loci 198, 331 and 554 deviated in two putative species (Supplementary Table S4). In most of these cases, there was evidence from MICRO-CHECKER for null alleles as the cause of the deviation (Supplementary Table S4).

Lumping individuals from different populations into a single group could lead to deviation from HWE and evidence for null alleles due to the Wahlund effect. To explore this possibility, we tested for deviation from HWE using GENEPOP and for evidence of null alleles using MICRO-CHECKER by testing each locus in each putative species at each sampling location separately (Supplementary Tables S4.1-S4.11). In the majority of cases (147 of 172 locus by species by locality combinations), genotypes did not deviate from HWE and there was no evidence of null alleles. These results suggest that all three putative species are regular outbreeders and that deviation from HWE was a consequence of having lumped individuals from different populations.

Interestingly, when we analysed deviation from HWE and sought evidence for null alleles by lumping individuals from different putative species into a single taxon, *B. lucorum s. lato*, either across all sampling localities (Supplementary Table S4) or for each sampling locality separately (Supplementary Tables S4.1-S4.11), loci were often out of HWE and showed evidence of null alleles (57 of 83 locus by location comparisons). These results provide a hint that the three putative species are differentiated in sympatry.

We tested for population genetic differentiation for each putative species separately across sampling localities for sampling site with  $n \geq 5$  individuals. For each putative species, differentiation across Ireland (Figure 2) was subtle, not significantly different from zero for

*B. cryptarum* (6 locations, global  $F_{ST} = 0.015$ ,  $P = 0.109$ ) but significant for *B. lucorum s. str.* (9 locations, global  $F_{ST} = 0.008$ ,  $P = 0.036$ ) and *B. magnus* (8 locations, global  $F_{ST} = 0.018$ ,  $P = 0.023$ ). For each putative species, Isolation by Distance was not significant (statistics and population pairwise  $F_{ST}$ , Supplementary Table S5), probably due to low statistical power (lack of sampling sites).

Differentiation between the three putative species was high (global  $F_{ST} = 0.268$ ,  $P < 0.001$ ); all three putative species pairs were significantly differentiated (*cryptarum* versus *lucorum s. str.*:  $F_{ST} > 0.205$ ,  $P < 0.001$ ; *cryptarum* versus *magnus*:  $F_{ST} > 0.219$ ;  $P < 0.001$ ; *cryptarum lucorum s. str.* versus *magnus*:  $F_{ST} > 0.322$ ,  $P < 0.001$ ). There was no suggestion in the  $F_{ST}$  values that *B. cryptarum* was closer to *B. magnus* than to *B. lucorum s. str.*, though *B. lucorum s. str.* was most distant to *B. magnus*. When we lumped the three putative species into one taxon, *B. lucorum s. lato*, differentiation across our 11 sampling sites in Ireland was insignificant ( $F_{ST} = 0.046$ , n.s.). These results suggest that the three putative species are genetically well differentiated.

Differences in allele frequencies of *B. lucorum s. str.* to the other two taxa were particularly evident at locus 327 (Figure 3). Allele frequencies also differed markedly in *B. magnus* compared to the other two putative species at locus 331 (Figure 3). Yet not one allele at any of the loci was both private (restricted to a putative species) and at a sufficiently high frequency within that species to allow it to be used to discriminate readily between species.

STRUCTURE analysis at  $K = 2$  (mean likelihood  $\ln \Pr(X|K) = -5668.6$ ; admixture ancestry model) revealed the mitochondrial lineage corresponding to *B. lucorum s. str.* to be well differentiated from *B. cryptarum* and *B. magnus* (Figure 4), which may reflect the closer

phylogenetic affinity of the latter pair of species than either of them to *B. lucorum s. str.* (Murray et al. 2008). The nuclear gene pools of the three putative species were clearly separated at  $K = 3$  (mean likelihood  $\text{Ln Pr}(X|K) = -5232.6$ ; Figure 4), with greater model support than for  $K = 1$  (mean likelihood  $\text{Ln Pr}(X|K) = -6880.2$ ) or  $K = 2$ . Results were qualitatively the same using STRUCTURE's non-admixture model (Supplementary Figure S1).

Multilocus microsatellite genotypes of one out of the 304 individuals did not concur with the COI mtDNA species delineation; an individual with a *magnus* mitochondrial haplotype was assigned to the *cryptarum* nuclear gene pool cluster (individual Ff53, Supplementary Table S6). Two additional individuals exhibited a major split in their nuclear gene pool assignments between two putative species (individuals Ff19 and Ff27:  $Q$  value  $\leq 0.5$ ; Supplementary Tables S6 and S7). STRUCTURE assignment of the other 301 individuals to their correct mitochondrial lineage was generally with high posterior probability ( $Q$  value  $> 0.93$ ); only 21 of the other 301 individuals (~8%) analysed exhibited a major assignment ( $Q$ ) value of  $< 0.9$ . These results lend weight to the hypothesis that the *lucorum* complex comprises three species, with good fit of multilocus nuclear genotypes to mitochondrial haplotypes and few exceptions.

Because STRUCTURE makes the strong assumption that genotypic data are in HWE, we repeated analyses by removing three loci that suggested marked deviation from HWE: loci 255, 331 and 198 (Supplementary Table S4). Results from STRUCTURE analyses with only four loci gave largely similar results to those with the entire dataset (Supplementary Figure S2).

Genotypes of the three mtDNA lineages each formed a separate cluster when mapped in multivariate space by PCoA, with only slight overlap at the edges of clusters (Figure 5). Seven of 304 individuals did not concur with COI mtDNA species delineation. These included the three individuals (Ff19, Ff27, Ff53) whose STRUCTURE assignment suggested their genotypes did not fit with other members of the same mitochondrial lineage (Supplementary Table S7).

Clustering genotypes by DAPC also revealed three discrete clusters that largely concurred with mitochondrial lineages (20 PCs extracted from the genotype data, Figure 6), providing additional support for the hypothesis that *B. cryptarum*, *B. lucorum s. str.* and *B. magnus* are discrete species. Five of 304 individuals were at the multivariate edge of mtDNA lineages. These also included the same three aberrant individuals (Ff19, Ff27, Ff53) highlighted by STRUCTURE and PCoA (Supplementary Table S7). The same five individuals were identified as outliers when 5, 10 or 40 PCs were extracted from the genotype data for DAPC (generating 7, 7 and 5 outlier individuals respectively).

Though STRUCTURE analysis suggested *B. cryptarum* and *B. magnus* are genetically closer to each other than either is to *B. lucorum s. str.* (Figure 4), PCoA and DAPC analyses did not support this view. Using the multivariate distance-based approaches, all three putative taxa were similarly differentiated (Figures 5 and 6).

## Discussion

Our multilocus nuclear (microsatellite) data of the *lucorum* complex of bumble bees collected in Ireland were grouped into three discrete multivariate clusters that corresponded well to the

three mitochondrial lineages formerly assigned to the putative species *B. cryptarum*, *B. lucorum s. str.* and *B. magnus*. Our data lend weight to the hypothesis, based on mtDNA COI partial sequences, that the *lucorum* complex comprises three morphologically cryptic but reproductively isolated species: *B. cryptarum*, *B. lucorum s. str.* and *B. magnus*.

Mallet (1995, 2007) has persuasively argued that, in the age of genetics and genomics, a robust species definition for sexually reproducing organisms is that, when in sympatry, they form discrete genotypic clusters. The multilocus genetic differentiation of the three *lucorum* complex mitochondrial lineages that we found in our STRUCTURE analyses at  $K = 3$  and using PCoA and DAPC is consistent with reciprocal monophyly of three species in sympatry.

Carstens et al. (2013) have argued that robust species delimitation should be based upon multiple, independent analyses. Here we employed three methods to differentiate nuclear gene pools of the *lucorum* complex, all of which were consistent in their identification of three discrete groups, with few outliers. STRUCTURE in particular makes the assumption of HWE, which, in our case, was violated at several loci when individuals from multiple locations were lumped together within a putative species. That STRUCTURE nevertheless identified the same outliers as PCoA and DAPC suggests it may be robust to given degree of violation of HWE in our dataset.

Though our nuclear microsatellite marker dataset suggests reproductive isolation between the three taxa of the *lucorum* complex, eight of 304 individuals possessed a multilocus genotype that did not concur with their mitochondrial haplotype. For three of these eight individuals, all three methods (STRUCTURE, PCoA and DAPC) placed them in a cluster different to that of other members of their mitochondrial lineage. One explanation for this incongruence is that

genotypic clusters of different species may overlap at their edges when based on a reduced number of loci; use of additional microsatellite loci or genome-wide markers may resolve this 'lack of data' issue (Lozier and Zayed 2016). Secondly, a low degree of hybridization between species may take place in the field. Individuals Ff19 and Ff27 could represent hybrids because their probability of assignment (STRUCTURE  $Q$  value) was intermediate between their mitochondrial lineage and that of another lineage. Artificial crosses between members of the *lucorum* complex are needed to support this suggestion. Thirdly, aberrant individuals might be a product of mitochondrial introgression; individual Ff53 could represent such a case, with a *magnus* mitochondrial haplotype confirmed by Sanger sequencing and a multilocus nuclear genotype assigned to *cryptarum*, *cryptarum/lucorum* and *cryptarum* by STRUCTURE, PCoA and DAPC respectively. It may be of significance that all three consistently aberrant individuals (Ff19, Ff27, Ff53) were collected at one site, Slieve Gullion, as workers (Table 1). We note, though, that the three discrete multilocus genotypic clusters we detected are unlikely to be maintained if hybridization or mtDNA introgression were common and widespread. Thus our second and third explanations seem unlikely to account for the mis-assigned individuals. Alternatively, if they do occur, they may not lead to fertile sexual descendants (queens and males).

If, as seems likely, our first explanation is correct, it suggests that considerable effort will be required for microsatellites to be used to separate among cryptic species or to detect hybridization. Within European bees, there are many putative cryptic species pairs or cryptic species complexes that share COI barcodes (Schmidt et al. 2015). Interspecific DNA sequence divergence at COI of the *lucorum* complex in Ireland is >2.3% (Murray et al. 2008), yet the 7 microsatellite loci we used to analyse 54-168 individuals per taxon were insufficient to resolve unambiguously all 304 individuals. For comparison, in the cryptic Neotropical



orchid bee sister species *Euglossa dilemma* and *Euglossa viridissima*, species-typical alleles at one locus have been found to differentiate between taxa (Eltz et al. 2011), though not unambiguously. Indeed, as the number of analysed individuals per taxon increases, so too is the likelihood of detecting greater allelic diversity, reducing the probability of finding private, species-diagnostic alleles at highly variable loci.

Lack of resolution in separating between reproductively isolated nuclear gene pools using microsatellites might be due to shared ancestral polymorphisms and homoplasy caused by high mutation rates at microsatellite loci (e.g. Schlötterer 1998). Sequence divergence of other cryptic species complexes of bee at COI is considerably less than 2.3%, suggesting a more recent common ancestor than that of the *lucorum* complex; for example, three members of the *Colletes succinctus* complex (species *hederae*, *halophilus* and *succinctus*) share the same COI barcode (Kuhlmann et al. 2007). For these and other closely related species, a larger sample size of sympatric individuals and, possibly more importantly, a larger number of nuclear loci may be needed to separate unambiguously among reproductively isolated nuclear gene pools (e.g. using deep sequencing via genome skimming, Cossia et al. 2016), calling into question the feasibility of using microsatellites for multilocus sequence typing so as to differentiate readily among cryptic species.

Species-specific pheromones may play an important role in mate-recognition, presumably decreasing the incidence of interspecific mating (Paterson 1985). *Bombus* male sex pheromones contain over 50 different volatile compounds derived from the labial glands that are used to scent-mark substrates and that are thought to attract unmated conspecific queens (Bergström et al. 1981; Ayasse et al. 2001). It has been previously demonstrated that males of *B. cryptarum*, *B. lucorum s. str.* and *B. magnus* differ in their cephalic secretions to the extent

that their multivariate chemical composition has been used to support specific classification of the three taxa (Bertsch 1997a; Bertsch et al. 2004, 2005; Pamilo et al. 1997). It is plausible that mate recognition based on male labial secretions plays a significant role as a prezygotic isolating mechanism in maintaining species boundaries between members of the *lucorum* complex.

The existence of three cryptic species within the *lucorum* complex of *Bombus* has several implications for conservation and management. In terms of conservation *per se*, members of the *lucorum* complex, like *B. terrestris*, are classified as of ‘least concern’ in Europe (Nieto et al. 2014). Nevertheless, it is highly likely that these cryptic species have been previously overlooked, and treatment of them as such should be borne in mind when addressing conservation status, particularly in light of the on-going declines in bumble bees in Europe (Goulson et al. 2005; Fitzpatrick et al. 2007; Rasmont et al. 2015). Indeed, in Ireland, *B. lucorum s. str.* is classified as ‘least concern’ whereas *B. cryptarum* and *B. magnus* are more cautiously and appropriately classified as ‘data deficient’ (Fitzpatrick et al. 2006). The three *lucorum* complex species may exhibit ecological specialization. With regard to altitude, *B. cryptarum* is has more of an upland distribution in Ireland (Murray et al. 2008) whereas *B. magnus* is considered a lowland species in Germany (von Hagen 2003). More recent analysis of their ecological associations has suggested that *B. cryptarum* and *B. magnus* prefer cooler sites in comparison to *B. lucorum* (Walters et al. 2010; Scriven et al. 2015). Phenological differences also exist; *B. cryptarum* is an early species that precedes *B. lucorum s. str.* and *B. magnus* (Bertsch 1997b; Bertsch et al. 2004), which may also play a role in preventing hybridization.

From a management perspective, it is necessary to be able to identify taxa correctly because bumble bees are important managed crop pollinators and colonies are increasingly being transplanted long distances to provide pollination services (Goulson 2003). This has become of particular conservation concern since bumble bee translocations have been implicated in colony declines, including through pathogen spillover, and in the replacement of native *Bombus* species (Inoue et al. 2008; Meeus et al. 2011; Schmid-Hempel et al. 2014). Indeed, in Asia, confusion remains over the identification of bumble bee species imported for crop pollination (Williams et al. 2012b). Importation of non-native bumble bee species not only brings risks associated with the introduction of a competitively superior congener and of a non-native's pathogens, there is also the risk of hybridization between native and introduced species. We note that inter-specific mating and hybridization can only be detected by using codominant nuclear markers such as microsatellites or SNPs; the former have been employed to demonstrate interspecific mating and hybrid inviability in crosses between native Japanese bumble bees and imported European *B. terrestris* (Kanbe et al. 2008; Tsuchida et al. 2010).

The utility of DNA barcoding versus morphological-based taxonomy in biodiversity inventorying has been hotly debated (Packer et al. 2009; Stahlhut et al. 2012; Gerth and Bleidorn 2013). Yet DNA barcodes continue to be used for species identification and even the characterization of new *Bombus* species (Williams et al. 2016). Our data vindicate their use for species identification within the *lucorum* complex.

**Acknowledgements** We thank friends and colleagues who helped to collect bumble bees across Ireland: D. Cookson, D. Dominoni, M. Kelly and S. Roos; Andreas Bertsch for use of his photographs, comments on this manuscript and encouragement to engage with the *lucorum* complex; and Robin Moritz for laboratory and intellectual support. We also thank

two anonymous reviewers and editor-in-chief as well as Shalene Jha and Christophe Praz for many insightful comments that helped improve the manuscript.

**Funding** This work was supported by a grant from the Higher Education Authority of Ireland as part of its North-South Research Programme for Peace and Reconciliation. L McKendrick thanks DARD for their financial support (a PhD stipend) and patience.

**Conflict of Interest** The authors declare that they have no conflict of interest.

## References

Ayasse M, Paxton RJ, Tengö J (2001) Mating behavior and chemical communication in the order Hymenoptera. *Ann Rev Entomol* 46:31-78.

Bergström G, Svensson BG, Appelgren M, Groth I (1981) Complexity of bumble bee marking pheromones: biochemical, ecological and systematic interpretations. pp.175-183 in: Hous PE, Clemen J-L (eds.) *Biosystematics of Social Insects* Vol. 19, Academic Press, London, UK.

Berkov A (2002) The impact of redefined species limits in *Palame* (Coleoptera: Cerambycidae: Lamiinae: Acanthocinini) on assessments of host, seasonal, and stratum specificity. *Biol J Linn Soc* 76:195-209.

Bertsch A (1997a) Abgrenzung der Hummel-Arten *Bombus cryptarum* und *B. lucorum* mittels männlicher Labialdrüsen-Sekrete und morphologischer Merkmale (Hymenoptera, Apidae). *Entomologia Generalis* 22:129-145.

- Bertsch A (1997b) Wieviele Arten der Untergattung *Terrestribombus* (Hymenoptera, Apidae) gibt es in Nordhessen; die Abgrenzung von *Bombus cryptarum* und *B. lucorum* mittels männlicher Labial-Drüsen-Sekrete und morphologischer Merkmale. Marburger Entomologische Publikationen 2:1-28.
- Bertsch A, Schweer H, Titze A (2004) Discrimination of the bumblebee species *Bombus lucorum*, *B. cryptarum* and *B. magnus* by morphological characters and male labial gland secretions. Beiträge zur Entomol 54:365-386.
- Bertsch A, Schweer H, Titze A, Tanaka H (2005) Male labial gland secretions and mitochondrial DNA markers support species status of *Bombus cryptarum* and *B. magnus* (Hymenoptera, Apidae). Insectes Soc 52:45-54.
- Bossert S (2015) Recognition and identification of bumblebee species in the *Bombus lucorum*-complex (Hymenoptera, Apidae) – A review and outlook. Dtsch Entomol Z 62:19-28.
- Cameron SA, Lozier JD, Strange JP, Koch JB, Cordes N, Solter LF, Griswold TL (2011) Patterns of widespread decline in North American bumble bees. Proc Nat Acad Sci USA 108:662–667.
- Carolan JC, Murray TE, Fitzpatrick Ú, Crossley J, Schmidt H, Cederberg B, McNally L, Paxton RJ, Williams PH, Brown MJF (2012) Colour patterns do not diagnose species: quantitative evaluation of a DNA barcoded cryptic bumblebee complex. PLoS ONE 7:e29251.
- Carstens BC, Pelletier TA, Reid NM, Satler JD (2013) How to fail at species delimitation. Mol Ecol 22:4369-4383.
- Coissac E, Hollingsworth PM, Lavergne S, Taberlet P (2016) From barcodes to genomes:

468 extending the concept of DNA barcoding. *Mol Ecol* 25:1423-1428.

469 Crandall KA, Binindamonds ORP, Mace GM, Wayne RK (2000) Considering evolutionary  
470 processes in conservation biology. *Trends Ecol Evol* 15:290-295.

471 Dieringer D, Schlötterer C (2003) MICROSATELLITE ANALYSER (MSA): a platform  
472 independent analysis tool for large microsatellite data sets. *Mol Ecol Notes* 3:167-169.

473 Eltz T, Frittsch F, Pech JR, Zimmermann Y, Ramírez SR, Quezada-Euan JJG, Bembé B  
474 (2011) Characterization of the orchid bee *Euglossa viridissima* (Apidae: Euglossini) and a  
475 novel cryptic sibling species, by morphological, chemical, and genetic characters. *Zool J*  
476 *Linn Soc* 163:1064-1076.

477 Fitzpatrick Ú, Murray TE, Byrne A, Paxton RJ, Brown MJF (2006) Regional red list of Irish  
478 bees. Dublin, Ireland: National Parks and Wildlife Service (Republic of Ireland) and  
479 Environment and Heritage Service (Northern Ireland). 1-38.

480 Fitzpatrick Ú, Murray TE, Paxton RJ, Breen J, Cotton D, Santorum V, Brown MJF (2007)  
481 Rarity and decline in bumblebees - a test of causes and correlates in the Irish fauna. *Biol*  
482 *Conserv* 136:185-194.

483 Gerth M, Bleidorn C (2013) A multilocus sequence typing (MLST) approach to diminish the  
484 problems that are associated with DNA barcoding: A reply to Stahlhut et al. (2012).  
485 *Systematics and Biodiversity* 11:1-3.

486 Gerth M, Gansauge M-T, Weigert A, Bleidorn C (2014) Phylogenomic analyses uncover  
487 origin and spread of the *Wolbachia* pandemic. *Nature Comm* 5:5117.

488 Gerth M, Geißler A, Bleidorn C (2011) *Wolbachia* infections in bees (Anthophila) and  
489 possible implications for DNA barcoding. *Systematics and Biodiversity* 9:319-327.

490 Gerth M, Röthe J, Bleidorn C (2013) Tracing horizontal *Wolbachia* movements among bees  
 491 (Anthophila): a combined approach using multilocus sequence typing data and host  
 492 phylogeny. *Mol Ecol* 22:6149-6162.

493 Goulson D (2003) Effects of introduced bees on native ecosystems. *Ann Rev Ecol Evol Syst*  
 494 34:1-26.

495 Goulson D (2009) Bumblebees. Behaviour, Ecology and Conservation. 2nd edn. Oxford  
 496 University Press, Oxford, UK.

497 Goulson D, Hanley ME, Darvill B, Ellis JS, Knight ME (2005) Causes of rarity in  
 498 bumblebees. *Biol Conserv* 122:1-8.

499 Hebert PDN, Cywinska A, Ball SL, deWaard JR (2003) Biological identifications through  
 500 DNA barcodes. *Proc Roy Soc Lond B* 270:313-321.

501 Hebert PDN, Penton EH, Burns JM, Janzen DH, Hallwachs W (2004) Ten species in one:  
 502 DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes*  
 503 *fulgerator*. *Proc Nat Acad Sci USA* 101:14812-14817.

504 Hines HM (2008) Historical biogeography, divergence times, and diversification patterns of  
 505 bumble bees (Hymenoptera: Apidae: *Bombus*). *Syst Biol* 57:58-75.

506 Inoue MN, Yokoyama J, Washitani I (2008) Displacement of Japanese native bumblebees by  
 507 the recently introduced *Bombus terrestris* (L.) (Hymenoptera: Apidae). *J Insect Cons*  
 508 12:135-146.

509 Jensen JL, Bohonak AJ, Kelley ST (2005) Isolation by distance, web service. *BMC Genet*  
 510 6:13.

511 Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers.

512        Bioinformatics 24:1403-1405.

513        Jombart T, Devillard S, Balloux F (2010) Discriminant analysis of principal components: a  
514        new method for the analysis of genetically structured populations. BMC Genet 11:94.

515        Kanbe Y, Okada I, Yoneda M, Goka K, Tsuchida K (2008) Interspecific mating of the  
516        introduced bumblebee *Bombus terrestris* and the native Japanese bumblebee *Bombus*  
517        *hypocrita sapporoensis results in inviable hybrids. Naturwiss 95:1003-1008.*

518        Kuhlmann M, Else GR, Dawson A, Quicke DLJ (2007) Molecular, biogeographical and  
519        phenological evidence for the existence of three western European sibling species in the  
520        *Colletes succinctus* group (Hymenoptera: Apidae). Organisms Divers Evol 7:155-165.

521        Lozier JD, Zayed A (2016) Bee conservation in the age of genomics. Conserv Genet this  
522        special issue.

523        Magnacca KN, Brown MJF (2010a) Mitochondrial heteroplasmy and DNA barcoding in  
524        Hawaiian *Hylaeus* (*Nesoprosopis*) bees (Hymenoptera: Colletidae). BMC Evol Biol  
525        10:174.

526        Magnacca KN, Brown MJF (2010b) Tissue segregation of mitochondrial haplotypes in  
527        heteroplasmic Hawaiian bees: implications for DNA barcoding. Mol Ecol Res 10:60-68.

528        Magnacca KN, Brown MJF (2012) DNA barcoding a regional fauna: Irish solitary bees. Mol  
529        Ecol Res 12:990-998.

530        Mallet J (1995) A species definition for the Modern Synthesis. Trends Ecol Evol 10:294-299.

531        Mallet J (2007) Species, concepts of. In: Levin SA et al., eds. Encyclopedia of Biodiversity.  
532        2nd ed. San Diego, California, USA: Academic Press. 427–440.

533        Meeus I, Brown MJF, De Graaf DC, Smagghe G (2011) Effects of invasive parasites on



534       bumble bee declines. *Conserv Biol* 25:662-671.

535   Murray TE, Fitzpatrick Ú, Brown MJF, Paxton RJ (2008) Cryptic species diversity in a  
536       widespread bumble bee complex revealed using mitochondrial DNA RFLPs. *Conserv*  
537       *Genet* 9:653-666.

538   Nieto A, Roberts SPM, Kemp J, Rasmont P, Kuhlmann M, García Criado M, Biesmeijer JC,  
539       Bogusch, Dathe HH, De la Rúa P, De Meulemeester T, Dehon M, Dewulf A, Ortiz-  
540       Sánchez FJ, Lhomme P, Pauly A, Potts SG, Praz C, Quaranta M, Radchenko VG,  
541       Scheuchl E, Smit J, Straka J, Terzo M, Tomozii B, Window J, Michez D (2014) European  
542       Red List of Bees. Publication Office of the European Commission: Luxembourg.

543   Packer L, Gibbs J, Sheffield CS, Hanner R (2009) DNA barcoding and the mediocrity of  
544       morphology. *Mol Ecol Res* 9:42-50.

545   Palsbøll PJ, Berube M, Allendorf FW (2007) Identification of management units using  
546       population genetic data. *Trends Ecol Evol* 22:11-16.

547   Pamilo P, Tengö J, Rasmont P, Pirhonen K, Pekkarinen A, Kaarnama E (1997) Pheromonal  
548       and enzyme genetic characteristics of the *Bombus lucorum* species complex in northern  
549       Europe. *Entomol Fennici* 14:187-194.

550   Paterson HE (1985) The recognition concept of species. In: Vrba ES (ed.) *Species and*  
551       *Speciation*. Transvaal Museum Monographs 4:21-29.

552   Paxton RJ, Thoren PA, Tengo J, Estoup A, Pamilo P (1996) Mating structure and nestmate  
553       relatedness in a communal bee, *Andrena jacobi* (Hymenoptera, Andrenidae), using  
554       microsatellites. *Mol Ecol* 5:511-519.

555   Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetic

556 software for teaching and research. Mol Ecol Notes 6:288-295.

557 Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using  
558 multilocus genotype data. Genetics 155:945-959.

559 R Core Team (2014). R: A language and environment for statistical computing. R Foundation  
560 for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>

561 Rasmont P (1984) Les bourdons du genre *Bombus* Latreille *sensu stricto* en Europe  
562 occidentale et centrale. Spixiana 7:135-160.

563 Rasmont P, Franzén M, Lecocq T, Harpke A, Roberts S, Biesmeijer JC, Castro L, Cederberg  
564 B, Dvorak L, Fitzpatrick Ú, Gonseth Y, Haubruge E, Mahé G, Manino A, Michez D,  
565 Neumayer J, Ødegaard F, Paukkunen J, Pawlikowski T, Potts S, Reemer M, Settele J,  
566 Straka J, Schweiger O (2015) Climatic risk and distribution atlas of European bumblebees.  
567 BioRisk 10:1-236.

568 Rasmont P, Scholl A, de Jonghe R, Obrecht E, Adamski A (1986) Identité et variabilité des  
569 mâles de bourdons du genre *Bombus* Latreille *sensu stricto* en Europe occidentale et  
570 centrale (Hymenoptera, Apidae, Bombinae). Rev Suisse Zool 93:661-682.

571 Raymond M, Rousset F (1995) GENEPOP (v. 1.2): Population genetic software for exact tests  
572 and ecumenicism. J Hered 86:248–249.

573 Riddle BR, Hafner DJ, Alexander LF, Jaeger JR (2000) Cryptic vicariance in the historical  
574 assembly of a Baja California Peninsular Desert biota. Proc Nat Acad Sci USA 97:14438-  
575 14443.

576 Ryder OA (1986) Species conservation and systematics: the dilemma of subspecies. Trends  
577 Ecol Evol 1:9-10.

578 Schlötterer C (1998) Microsatellites. In: Molecular Genetic Analysis of Populations. A  
579 Practical Approach (ed. Hoelzel AR), pp. 237-261. Oxford University Press, Oxford, UK.

580 Schmid-Hempel R, Eckhardt M, Goulson D, Heinzmann D, Lange C, Plischuk S, Escudero  
581 LR, Salathé R, Scriven JJ, Schmid-Hempel P (2014) The invasion of southern South  
582 America by imported bumblebees and associated parasites. *J Anim Ecol* 83:823-837.

583 Scriven JJ, Woodall LC, Tinsley MC, Knight ME, Williams PH, Carolan JC, Brown MJF,  
584 Goulson D (2015) Revealing the hidden niches of cryptic bumblebees in Great Britain:  
585 implications for conservation. *Biol Conserv* 182:126-133.

586 Schmidt S, Schmid-Egger C, Morinière J, Haszprunar G, Hebert PDN (2015) DNA barcoding  
587 largely supports 250 years of classical taxonomy: identifications for Central European  
588 bees (Hymenoptera, Apoidea *partim*). *Mol Ecol Res* 15:985-1000.

589 Sheffield CS, Hebert PDN, Kevan PG, Packer L (2009) DNA barcoding a regional bee  
590 (Hymenoptera: Apoidea) fauna and its potential for ecological studies. *Mol Ecol Res*  
591 9:196-207.

592 Stahlhut JK, Gibbs J, Sheffield CS, Alex Smith M, Packer L (2012) *Wolbachia*  
593 (Rickettsiales) infections and bee (Apoidea) barcoding: a response to Gerth et al.  
594 *Systematics and Biodiversity* 10: 395-401.

595 Stolle E, Wilfert L, Schmid-Hempel L, Schmid-Hempel P, Kube M, Reinhardt R, Moritz  
596 RFA (2011) A second generation genetic map of the bumblebee *Bombus terrestris*  
597 (Linnaeus, 1758) reveals slow genome and chromosome evolution in the Apidae. *BMC*  
598 *Genomics* 12:48.

599 Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular  
600 evolutionary genetics analysis using maximum likelihood, evolutionary distance, and

601 maximum parsimony methods. Mol Biol Evol 28:2731-2739.

602 Tsuchida K, Ito Kondo N, Inoue MN, Goka K (2010) Reproductive disturbance risks to  
 603 indigenous Japanese bumblebees from introduced *Bombus terrestris*. Appl Entomol Zool  
 604 45:49-58.

605 Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER:  
 606 software for identifying and correcting genotyping errors in microsatellite data. Mol Ecol  
 607 Notes 4:535-538.

608 Vesterlund SR, Sorvari J, Vasemägi A (2014) Molecular identification of cryptic bumblebee  
 609 species from degraded samples using PCR–RFLP approach. Mol Ecol Res 14:122-126.

610 von Hagen E (2003) Hummeln: Bestimmen, Ansiedeln, Vermehren, Schützen. Fauna-Verlag,  
 611 Nottuln, Germany.

612 Walsh PS, Metzger DA, Higuchi R (1991) Chelex 100 as a medium for simple extraction of  
 613 DNA for PCR-based typing from forensic material. Biotechniques 10:506-513.

614 Waters J, Darvill B, Lye GC, Goulson D (2011) Niche differentiation of a cryptic bumblebee  
 615 complex in the Western Isles of Scotland. Insect Conserv Div 4:46-52.

616 Williams PH (1998) An annotated checklist of bumble bees with an analysis of patterns of  
 617 description (Hymenoptera: Apidae, Bombini). Bull Natural History Mus Lond (Entomol)  
 618 67:79-152.

619 Williams PH, Cameron SA, Hines HM, Cederberg B, Rasmont P (2008) A simplified  
 620 subgeneric classification of the bumblebees (genus *Bombus*). Apidol 39:46-74.

621 Williams PH, Brown MJF, Carolan JC, An J, Goulson D, Aytekin AM, Best LR, Byvaltsev  
 622 AM, Cederberg B, Dawson R, Huang J, Ito M, Monfared A, Raina RH, Schmid-Hempel

623 P, Sheffield CS, Šima P, Xie Z (2012a) Unveiling cryptic species of the bumblebee  
 624 subgenus *Bombus s. str.* worldwide with COI barcodes (Hymenoptera: Apidae).  
 625 Systematics and Biodiversity 10:21-56.

626 Williams PH, An J, Brown MJF, Carolan JC, Goulson D, Huang J, Ito M (2012b) Cryptic  
 627 bumblebee species: consequences for conservation and the trade in greenhouse pollinators.  
 628 PLoS ONE 7:e32992.

629 Williams PH, Cannings SG, Sheffield CS (2016) Cryptic subarctic diversity: a new  
 630 bumblebee species from the Yukon and Alaska (Hymenoptera: Apidae). J Nat Hist:1-13.  
 631 DOI: 10.1080/00222933.2016.1214294

## Figure legends

**Fig. 1** Queens of the four Irish members of the *Bombus s. str.* group, namely *B. terrestris* and the three members of the *lucorum* complex: *B. cryptarum*, *B. lucorum s. str.* and *B. magnus* (photo credit: Andreas Bertsch 2004)

**Fig. 2** Sample sites (numbers correspond to those in Table 1) and numbers of individuals genotyped at seven microsatellite loci of the *lucorum* complex of bumble bees. Species designation was based on mtDNA COI RFLPs and updated by Sanger sequencing

**Fig. 3** Bubble plots of allele frequencies at seven microsatellite loci of 304 individuals of the three putative species of the *lucorum* complex of bumble bees from 11 sites in Ireland; species designation was based on mtDNA COI haplotypes and bubble diameters reflect allele frequencies which, within a species (column), sum to one

**Fig. 4** Barplot of STRUCTURE (using the admixture ancestry model) output showing percentage assignment of individuals of the three *lucorum* complex species of bumble bees genotyped at seven microsatellite loci to a given putative species for  $K = 2$  and  $K = 3$  (species designation based on mtDNA COI haplotypes)

**Fig. 5** PCoA of the multilocus microsatellite genotypes of 304 *lucorum* complex bumble bees from Ireland; each of the three mtDNA lineages is coded by a different shading; the three individuals (Ff19, Ff27, Ff53) with low multilocus group membership (low STRUCTURE  $Q$  value) to others of the same mtDNA lineage are labelled and circled in grey

657 **Fig. 6** DAPC of the multilocus microsatellite genotypes of 304 *lucorum* complex bumble  
658 bees from Ireland; each of the three mtDNA lineages is coded by a different shading; the  
659 three individuals (Ff19, Ff27, Ff53) with low multilocus group membership (low STRUCTURE  
660  $Q$  value and DAPC assignment to a different lineage) to others of the same mtDNA lineage  
661 are labelled and circled in grey  
662

663 **Table 1** Sample sites in Ireland and numbers of each species (as defined by mitochondrial lineage) of 304 *Bombus lucorum sensu lato* (*B.*  
664 *cryptarum*, *B. lucorum s. str.* and *B. magnus*) collected and genotyped at 5-7 microsatellite loci (see Table 1 of Murray et al. (2008) for dates  
665 of specimen collection); specimens were queens except where given in parentheses as workers (w)

| No           | Location                 | Latitude (N) | Longitude (W) | <i>B. cryptarum</i> | <i>B. lucorum s. str.</i> | <i>B. magnus</i> |
|--------------|--------------------------|--------------|---------------|---------------------|---------------------------|------------------|
| 1            | Dublin City, Co. Dublin  | 53°20'22"    | 06°13'38"     | 11                  | 28                        | 0                |
| 2            | Clara, Co. Wicklow       | 52°58'07"    | 06°15'59"     | 2                   | 13                        | 17               |
| 3            | Glenasmole, Co Dublin    | 53°13'54"    | 06°20'52"     | 3                   | 12                        | 8                |
| 4            | Glencree, Co. Wicklow    | 53°11'41"    | 06°18'22"     | 9                   | 4                         | 7                |
| 5            | Kippure, Co. Wicklow     | 53°10'47"    | 06°18'18"     | 6                   | 3                         | 11               |
| 6            | Powerscourt, Co. Wicklow | 53°09'59"    | 06°14'55"     | 5                   | 6                         | 8                |
| 7            | Killarney, Co. Kerry     | 52°03'48"    | 09°29'55"     | 0                   | 19                        | 12               |
| 8            | Belfast City, Co. Antrim | 54°34'13"    | 05°55'09"     | 6 (w=5)             | 18 (w=12)                 | 0                |
| 9            | Cork City, Co. Cork      | 51°53'54"    | 08°25'29"     | 1                   | 40                        | 0                |
| 10           | Slieve Gullion, Co. Down | 54°06'47"    | 06°24'55"     | 7 (all w)           | 18 (all w)                | 14 (all w)       |
| 11           | Benbulbin, Co. Sligo     | 54°18'49"    | 08°23'13"     | 4 (all w)           | 7 (all w)                 | 5 (all w)        |
| <b>Total</b> |                          |              |               | <b>54</b>           | <b>168</b>                | <b>82</b>        |



## Supplementary files

**Table S1** Microsatellite primers developed for *Bombus terrestris* by Stolle et al. (2011; see Table S1 of Stolle et al. 2011) and employed in this study to genotype the *lucorum* complex of three putative species: *B. cryptarum*, *B. lucorum s. str.* and *B. magnus*

**Table S2.** Details of the eight individuals of the *lucorum* complex whose RFLP-based mt haplotype did not concur with nuclear (microsatellite) genotypes, as defined by STRUCTURE  $Q$  value (Supplementary Table S6) or visually by PCoA (Figure 5). Code: the unique identifier to the specimen; Location: site of collection; RFLP ID: haplotype designation by RFLP (from Murray et al. 2008); PCoA ID: microsatellite genetic cluster membership (see Figure 5); STRUCTURE  $Q$  ID: microsatellite genetic cluster membership by STRUCTURE  $Q$  value (see Supplementary Table S6); COI-BLAST: COI barcode DNA sequence identity by BLASTn search against entire NCBI nucleotide database; GenBank Accession Number of sequence with 100% identity to Irish sample

**Table S3** Linkage disequilibrium for seven microsatellite loci in the three putative species of the *lucorum* complex of bumble bees collected at 11 sites in Ireland; columns represent respectively the pairwise combination of loci, probability of linkage disequilibrium (P) and standard error of that P value (SE), calculated in GENPOP

**Table S4** Deviation from Hardy-Weinberg equilibrium (HWE) for seven microsatellite loci in the three putative species of the *lucorum* complex of bumble bees collected from 11 sites in Ireland, including the observed and expected heterozygosity ( $H_o$  and  $H_{exp}$  respectively), the inbreeding coefficient,  $F_{IS}$ , the probability of deviation from HWE as calculated in GENEPOP

v. 4.2 (<http://genepop.curtin.edu.au>), and evidence for null alleles in MICRO-CHECKER; also given are  $H_o$ ,  $H_{exp}$ ,  $F_{IS}$ , and the probability of deviation from HWE for all *B. lucorum s. l.* collected from site 1 and combined into one group. **Tables S4.1-S4.11** give the same details for sampling sites 1-11 respectively.

**Table S5** Semi-matrix of pairwise estimates of  $F_{ST}$  across sampling locations (a) of *B. cryptarum* (n = 6 locations), (b) *B. lucorum s. str.* (n = 9 locations) and (c) *B. magnus* (n = 8 locations), for which the number of individuals genotyped per location  $\geq 5$  (location notation follows that given in Figure 1); no pairwise values are significantly different from zero ( $P < 0.05$ ) using MSA v. 4.05 (Dieringer and Schlötterer 2003)

**Table S6** Mitochondrial lineage and multilocus probability of group membership (STRUCTURE  $Q$  value; admixture model) at seven microsatellite loci in the three putative species of the *lucorum* complex of bumble bees collected at 11 sites in Ireland (see Table 1 for site names and location); the one individual with aberrant correspondence between mtDNA lineage (mt sp.) and nuclear genotype (nuc cryp, nuc luc, nuc mag for *B. cryptarum*, *B. lucorum s. str.* and *B. magnus*, respectively) is highlighted in blue; the two individuals with borderline membership to their mtDNA lineage (STRUCTURE  $Q$  value  $\leq 0.5$ ) are highlighted in green; the 21 individuals with non-aberrant but low probability (STRUCTURE  $Q$  value  $> 0.5$  and  $< 0.9$ ) of membership to their mtDNA lineage are highlighted in yellow


**Table S7** Details of the eight individuals of the *lucorum* complex whose updated mt haplotype did not concur with nuclear (microsatellite) genotypes, as defined by STRUCTURE  $Q$  value (Supplementary Table S6) or visually by PCoA (Figure 5) or by DAPC assignment (Figure 6). Code: the unique identifier to the specimen; Location: site of collection;

Haplotype ID: haplotype designation by RFLP (from Murray et al. 2008) or Sanger sequencing; STRUCTURE  $Q$  ID: microsatellite genetic cluster membership by STRUCTURE  $Q$  value (see Supplementary Table S6); PCoA ID: microsatellite genetic cluster membership (see Figure 5); DAPC ID: microsatellite genetic cluster membership (see Figure 6); comment: possible cause for lack of mt-nuclear concurrence

**Fig. S1** Barplot of STRUCTURE (using the non-admixture model) output showing percentage assignment of individuals of the three *lucorum* complex species of bumble bees genotyped at seven microsatellite loci to a given putative species for  $K = 2$  and  $K = 3$  (species designation based on mtDNA COI haplotypes)

**Fig. S2** Barplot of STRUCTURE (using admixture model) output showing percentage assignment of individuals of the three *lucorum* complex species of bumble bees genotyped at only four microsatellite loci to a given putative species for  $K = 3$  (species designation based on mtDNA COI haplotypes). STRUCTURE criteria were the same as in the ms for all seven loci (modelling without priors, and with the admixture ancestry model, burn-in of 50,000 followed by a further 500,000 iterations). The data fit a model with  $K=3$  better (mean likelihood  $\ln \Pr(X|K) = -3206$ ) than with  $K=2$  ( $\ln \Pr(X|K) = -3343$ ) or  $K=1$  ( $\ln \Pr(X|K) = -3997$ ). However, STRUCTURE gave two optimal results with  $K=3$  for the reduced dataset of four loci, (a) a poorer fit (probably local) optimum in which ‘cryptarum’ and ‘magnus’ formed one group whilst ‘lucorum’ was split into two groups (6 of 20 runs,  $\ln \Pr(X|K) = -3340$ ) and (b) a better fit (probably global) optimum as in our original results presented in Fig 4 (14 of 20 runs,  $\ln \Pr(X|K) = -3148$ ). All 20 runs converged.

Figure 1





**Figure 2**

Figure 3

Figure 3





**Figure 4**

Figure 5

Principal Coordinates Analysis (PCoA)





**Figure 6****Discriminant Analysis of Principal Components (DAPC)**